



How to Use an Auvi-Q for Anaphylaxis

Key points to remember:



- **Make sure you know how to use your child's auto-injector** so you are prepared in case of an emergency. Talk to your child's allergist or allergy nurse if you have questions about how to use the auto-injector.
- **If your child has symptoms of anaphylaxis, use the auto-injector and call 911 right away.**

What is anaphylaxis?

- Anaphylaxis is a very bad allergic reaction that can affect the whole body. **It is a medical emergency.**
- Anaphylaxis can happen a few seconds to a few hours after being exposed to the allergen (the substance that causes the reaction).
 - This allergic reaction most often happens within the first 30 minutes after being exposed to the allergen.
- Anaphylaxis can be very serious and sometimes deadly.
- Ask your allergist for a written emergency action plan.
- If possible, staying away from the allergen is best.
- Wearing a medic alert bracelet is also a good idea.

What are the symptoms?

Symptoms may include:



If your child has symptoms of anaphylaxis, use the epinephrine auto-injector and call 911 right away.

What is an epinephrine auto-injector?

- An epinephrine auto-injector is a device that gives 1 dose of epinephrine, also called adrenaline.
- It is important to talk with your health care provider about when to use it.

What should I know about using epinephrine auto-injectors?

- Keep enough epinephrine auto-injectors with you so there are always at least 2 with you or your child.
- Do not keep the epinephrine auto-injector in places where it is very hot or cold, like a car or glove compartment. It may cause the device to not work.

What is the medication in the auto-injector?

The medicine is called epinephrine. It works quickly to reverse symptoms of anaphylaxis/allergic reaction. It increases the heart rate and blood pressure, makes breathing easier and reduces the swelling around the lips, face and airway. Side effects may include:

- Fast/fluttery heartbeat, sweating, anxiety
- Dizziness
- Headache

Contact us

- **Monday–Friday, 8 a.m.–4:30 p.m.:** Call the Allergy Program at 617-355-6117.
- **After 4:30 p.m., on weekends and holidays:** Call 617-355-6369 to speak with the allergist on call.

This Family Education Sheet is available in Arabic and Spanish.

How do I use the Auvi-Q?




1 Remove Auvi-Q from the case.




2 Remove red safety guard.



3 Place against outer middle thigh.



4 Press firmly until you hear a "click" and "hiss" sound. **Hold in place for 2 seconds.**



5 Call 911.



6 Use 2nd Auvi-Q if symptoms are not better within 5 minutes.